

Calculating Correlation

by Sophia

	WHAT'S COVERED
Th fo	nis tutorial will introduce correlation and correlation coefficient. Our discussion breaks down as Ilows:
	1. Correlation
	a. Calculating Using Formula
	b. Calculating Using Excel Function

2. Scatter Plots with Different Correlation Coefficients

1. Correlation

When first describing scatter plots, you learned about their form, direction, and strength.

- Form assessed the linearity
- Direction says whether the data points tend to move in a positive or negative direction
- Strength shows how well they follow that form.

When the form is linear, we can use a number called **correlation**. It measures the strength and direction of a linear relationship. The direction will be easy to spot. It will be a positive number if there's a positive association, and a negative number if there's a negative association. The numerical quantity will measure strength.

😭 BIG IDEA

In terms of direction, what correlation would do is this: if the explanatory and response variables rise together, that's going to be called the positive direction. If one falls as the other increases, that's going to be called a negative correlation.

The correlation is measured using a numerical value known as the **correlation coefficient**. The correlation coefficient is a variable called "r" and is unit-less. It is expressed as a number between negative 1 and positive 1 and indicates the strength of the linear association.

Numbers that are close to negative 1 or positive 1 are associated with a strong association between the two variables--a 1 indicating a strong positive association, and a negative 1 indicating a strong negative association. Numbers near zero represent almost no linear relationship.





You can use the chart above to help you to understand the value of a correlation. Numbers between 0.8 and 1 are considered to have a strong correlation; between 0.5 and 0.8 a moderate correlation; and between 0 and 0.5 a very weak correlation. The same exists between negative 1 and 0.



Correlation

The strength and direction of a linear association between two quantitative variables.

Correlation Coefficient

The numerical value between -1 and +1 that measures the correlation between two quantitative variables.

1a. Calculating Using Formula

Correlation is essentially the average of the products of the z-scores for the x's and the y's. The z- scores are the values of x minus the means of x divided by the standard deviation of x. It's the same thing for y.

FORMULA TO KNOW

Correlation

$$r = \frac{1}{n-1} \sum z_x \cdot z_y = \frac{1}{n-1} \sum (\frac{x-\overline{x}}{s_x})(\frac{y-\overline{y}}{s_y})$$

 \Rightarrow EXAMPLE These are destinations that you could go to from the city of Minneapolis-Saint Paul, with the distances away from Minneapolis and the airfare to fly to any of these places.

Destination	Miles	Airfare
Kansas City	460	379
Los Angelas	1,870	377
Milwaukee	338	158
New York City	1,167	283
Philadelphia	1,141	323

Std.

Step 1: Calculate z-scores of the x variable. In this situation, miles is x, or the explanatory variable, as miles are believed to cause airfare to rise. This makes airfare the response variable, y. Take the given miles and airfare and convert both of them into z-scores.

To do this, you need the mean and the standard deviation. Recall from Unit 3 that you can use Microsoft Excel to easily find these values. For the mean, use the function "=AVERAGE", and for the standard deviation, use the function "=STDEV.S". When using these functions in Excel, you just need to highlight each column that you are finding the mean and standard deviation for. We will do this for both Miles and Airfare.

Miles	Airfare	Destination	Miles	Airfare	
460	379	Kansas City	460	379	
1870	377	Los Angeles	1870	377	
338	158	Milwaukee	338	158	
1167	283	New York City	1167	283	
1141	323	Philadelphia	1141	323	
=AVERAGE(C4:C8)	Mean	995.2	=AVERAGE(D4:D8)
		Std. Dev			
	Miles 460 1870 338 1167 1141 =AVERAGE(Miles Airfare 460 379 1870 377 338 158 1167 283 1141 323 =AVERAGE(C4:C8)	Miles Airfare Destination 460 379 Kansas City 1870 377 Los Angeles 338 158 Milwaukee 1167 283 New York City 1141 323 Philadelphia =AVERAGE(C4:C8) Mean Std. Dev	Miles Airfare Destination Miles 460 379 Kansas City 460 1870 377 Los Angeles 1870 338 158 Milwaukee 338 1167 283 New York City 1167 1141 323 Philadelphia 1141 =	Miles Airfare 460 379 1870 377 338 158 1167 283 1141 323 Philadelphia 1141 328 Mean 995.2 Std. Dev

Destination	Miles	Airfare	Destination	Miles	Airfare	
Kansas City	460	379	Kansas City	460	379	
Los Angeles	1870	377	Los Angeles	1870	377	
Milwaukee	338	158	Milwaukee	338	158	
New York City	1167	283	New York City	1167	283	
Philadelphia	1141	323	Philadelphia	1141	323	
Mean	995.2	304	Mean	995.2	304	
Std. Dev	=STDEV.S(C4	4:C8)	Std. Dev	619.35426	=STDEV.S(D	4:D9)

			~
Destination	Miles	Airfare	
Kansas City	460	379	
Los Angeles	1870	377	
Milwaukee	338	158	
New York City	1167	283	
Philadelphia	1141	323	
Mean	995.2	304	
Std. Dev	619.35426	90.928543	

Next, to calculate the z-score, subtract the mean from each value and divide by the standard deviation. For example, using the first value in Miles, take 460 minus the mean, 995.2, and divide by the standard deviation, 619.35. This gives us a -0.864.

		<u>460 -</u> 61	<u>995.2</u> 9.35
Destination	Miles	Airfare	z _x
Kansas City	460	379	-0.864126
Los Angeles	1870	377	1.412439
Milwaukee	338	158	-1.061105
New York City	1167	283	0.277386
Philadelphia	1141	323	0.235406
Mean	995.2	304	
Std. Dev	619.35426	90.928543	

Do the same thing for the 1870 miles to Los Angeles, and all of the other cities.

Step 2: Repeat this process and calculate the z-scores for the y values. In this scenario, the response

variables are the airfare values. Starting with Kansas City, 379 minus 304 divided by 90.93 gives us 0.825.

			379	- <u>304</u> 93
Destination	Miles	Airfare	z _x	Zy Zy
Kansas City	460	379	-0.864126	0.824824
Los Angeles	1870	377	1.412439	0.802828
Milwaukee	338	158	-1.061105	-1.605656
New York City	1167	283	0.277386	-0.230951
Philadelphia	1141	323	0.235406	0.208955
				sum
Mean	995.2	304		
Std. Dev	619.35426	90.928543		

Do the same thing with all the rest of the airfare.

Step 3: Multiply the corresponding z-scores and add. Starting with -0.864 and 0.825, multiply the

corresponding z-scores for the x and y variables, all down the rows, then add them up.

Destination	Miles	Airfare	z _x	z _y	z _x * z _y
Kansas City	460	379	-0.864126	0.824824	-0.712751
Los Angeles	1870	377	1.412439	0.802828	1.1339457
Milwaukee	338	158	-1.061105	-1.605656	1.7037703
New York City	1167	283	0.277386	-0.230951	-0.064062
Philadelphia	1141	323	0.235406	0.208955	0.0491894
				sum	2.110092
Mean	995.2	304			
Std. Dev	619.35426	90.928543			

The sum here ends up being positive 2.11. We can substitute this value into the correlation formula.

$$r = \frac{1}{n-1} \sum z_x \cdot z_y$$
$$r = \frac{1}{n-1} \sum 2.11$$

Step 4: Finally, divide by the number of observations minus 1. There are five observations, so the denominator will be 5-1.

$$r = \frac{1}{n-1}(2.11)$$
$$r = \frac{1}{5-1}(2.11)$$
$$r = \frac{1}{4}(2.11)$$
$$r = 0.527$$

Dividing by four yields a correlation of 0.527. This value tells us that the correlation between airfare and miles is a positive relationship but fairly weak association. We can also see this from the scatter plot:



1b. Calculating Using Excel Function

This is a *very* cumbersome process to go through, and the correlation coefficient is almost always found using technology. In Excel, once we have the basic information for miles and airfare listed, all you have to do is type in the command "=CORREL", which is short for correlation. Select all the things believed to be the x's, and all of the things we believe to be the y's. Close the parentheses and hit "Enter."

Miles	Airfare
460	379
1870	377
338	158
1167	283
1141	323
=correl(C3	:C7,D3:D7)
	Miles 460 1870 338 1167 1141 =correl(C3

Sure enough, it gives you the 0.527 that you got before.

2. Scatter Plots with Different Correlation Coefficients

Let's explore some scatter plots with different correlation coefficients.

Graph	Correlation	Explanation
	r = -0.99	The data points are in a negative direction, so the correlation is a negative number. It is also nearly linear, so its correlation is negative 0.99, which is very close to negative 1. This graph shows a very strong, negative association.
	r = -0.5	The data points are in a negative direction, so the correlation is a negative number. However, the data is fairly spread out, so the strength is not terribly strong. This graph shows a weak to moderate, negative association.
	r = 0	The data points have a cloudy association, so it is neither positive nor negative. There is also no linear association between the two variables, so the correlation is zero.
	r = 0.7	The data points show an upward association, so the correlation is a positive number. Although it is linear, the data points are not very clustered, but still close enough to show a fairly moderate to strong association.
	r = 0.9	The data points are in a positive direction, so the correlation is a positive number. It is also very linear, so its correlation will be closer to 1. This graph shows a strong, positive association.

	r = 0.3	Even though the points are spread out, a positive association is visible. However, since they are not closely clustered, the data points will show a weaker strength.
--	---------	--

The correlation only relates the linear relationship between two *quantitative* variables. As a caution, you're going to hear the word correlation thrown around a lot in everyday speech; however, there are often very common errors made when comparing two different variables. It is always important to make sure the two variables being measured are quantitative.

🕸 THINK ABOUT IT

Can you spot the errors in the following statements?

"There is a strong correlation between Type 2 diabetes, physical inactivity, and obesity."

Although it's possible that they are related, you can't use the word correlation. The first error here is that three variables are being compared, and correlation only compares two variables. Also, Type 2 diabetes is categorical--either you have it, or you don't. Physical inactivity could be quantitative, but it's not obviously quantitative. Obesity is certainly categorical.

"There is a strong correlation between IQ and religious affiliation."

Although IQ is quantitative, religious affiliation is categorical. You can't calculate the correlation between the two.

SUMMARY

Correlation measures the strength and direction of a linear relationship between two variables on a scatter plot. Strong associations have correlation coefficients near positive 1 or negative 1. Scatterplots with weak correlation coefficients are values near zero. Almost always, you can find it using technology, such as a calculator, an Internet Applet, or a spreadsheet. Because of the way that r is calculated, where you're multiplying z-scores, it doesn't matter which variable is called the explanatory and which is the response.

Good luck!

TERMS TO KNOW

Correlation

The strength and direction of a linear association between two quantitative variables.

Correlation coefficient (r)

The numerical value between -1 and +1 that measures the correlation between two quantitative variables.

FORMULAS TO KNOW

Correlation

$$r = \frac{1}{n-1} \sum z_x \cdot z_y = \frac{1}{n-1} \sum \left(\frac{x-\overline{x}}{s_x} \right) \left(\frac{y-\overline{y}}{s_y} \right)$$