

# Calculating Standard Deviation

by Sophia



## WHAT'S COVERED

This tutorial will explain the concept of standard deviation. Our discussion breaks down as follows:

1. Standard Deviation
2. Calculating Standard Deviation
3. Interpreting Standard Deviation
4. Using Technology

## 1. Standard Deviation

**Standard deviation** is a measure of variation that we use quite often in statistics. Standard deviation measures spread. You will interpret the standard deviation as the typical amount that you would expect data to be within the mean.

The full name for standard deviation is "standard deviation from the mean." If you break that down, "standard" just means it's typical, "deviation" means that you expect it to be off from the mean, just by chance. So "standard deviation from the mean" states that the data will be away from the mean.

Here is what the formula to find the standard deviation of a sample looks like:



### FORMULA TO KNOW

#### Standard Deviation of a Sample

$$s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$$

$n$  = number of data points

$x_i$  = each of the values in the data set

$\bar{x}$  = mean of the data set

As complicated as the formula looks, it's actually the one preferred most, provided that the distribution that you're looking for is roughly symmetric and doesn't have outliers. If the data is not symmetric or has outliers,

you will use a different measure of spread, the interquartile range.



#### TERM TO KNOW

#### Standard Deviation

A typical amount by which we would expect a data point to differ from the mean. Typically, about half to two-thirds of the data points fall within one standard deviation of the mean.

## 2. Calculating Standard Deviation

When calculating the formula for standard deviation, you follow these steps:



#### STEP BY STEP

**Step 1:** Subtract the mean from each value.

**Step 2:** Square those values, resulting in  $(x \text{ minus mean})^2$ .

**Step 3:** Use the sigma notation, which is the same as summation notation, to add these values.

**Step 4:** Divide that sum by  $(n \text{ minus } 1)$ .

**Step 5:** Take the square root.

$$s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$$

$n$  = number of data points

$x_i$  = each of the values in the data set

$\bar{x}$  = mean of the data set

↪ **EXAMPLE** These are the heights of the Chicago Bulls basketball team.

Height of Chicago Bulls Players	
Omer Asik	84
Carlos Boozer	81
Ronnie Brewer	79
Jimmy Butler	79
Luol Deng	81
Taj Gibson	81
Richard Hamilton	79
Mike James	74
Kyle Korver	79

John Lucas III	71
Joakim Noah	83
Derrick Rose	75
Brian Scalabrine	81
Marquis Teague	74
C.J. Watson	74

When considering standard deviation, each of these items is a  $X_i$  ( $X$  sub  $i$ ) in the original formula. Let's follow the five steps to calculate the standard deviation:

### STEP BY STEP

**Step 1:** The first step is to subtract the mean, which in this case, means you have to first calculate the mean. A thorough explanation of how to calculate the mean is covered in another tutorial, so for today, know that the mean is around 78.33 inches. Therefore, subtract 78.33 from each of these height values.

84 minus 78.33 is 5.67, 81 minus 78.33 is 2.67, etc.

$X$	$X - \bar{X}$
84	5.67
81	2.67
79	0.67
79	0.67
81	2.67
81	2.67
79	0.67
74	-4.33
79	0.67
71	-7.33
83	4.67
75	-3.33
81	2.67
74	-4.33
74	-4.33

**Step 2:** Square those values, resulting in  $(x - \bar{x})^2$ .  $5.67^2$  is 32.15,  $2.67^2$  is 7.13, etc.

$x$	$x - \bar{x}$	$(x - \bar{x})^2$
84	5.67	32.15
81	2.67	7.13
79	0.67	0.45
79	0.67	0.45
81	2.67	7.13
81	2.67	7.13
79	0.67	0.45
74	-4.33	18.75
79	0.67	0.45
71	-7.33	53.73
83	4.67	21.81
75	-3.33	11.09
81	2.67	7.13
74	-4.33	18.75
74	-4.33	18.75

**Step 3:** Use this sigma notation, which is the same as summation notation, to add these values up. They sum up to 205.35.

$x$	$x - \bar{x}$	$(x - \bar{x})^2$
84	5.67	32.15
81	2.67	7.13
79	0.67	0.45
79	0.67	0.45
81	2.67	7.13
81	2.67	7.13
79	0.67	0.45
74	-4.33	18.75

79	0.67	0.45
71	-7.33	53.73
83	4.67	21.81
75	-3.33	11.09
81	2.67	7.13
74	-4.33	18.75
74	-4.33	18.75
Sum		205.35

**Step 4:** Divide that sum by n minus 1. In this case, n is 15 because there were 15 players; therefore n minus 1 equals 14. Dividing our sum by 14 equals 14.67.

$$\frac{1}{15-1}(205.35) = \frac{1}{14}(205.35) = 14.67$$



HINT

If you stopped here, this 14.67 number would measure another kind of variation called "**variance**." Variance is equivalent to the square of the standard deviation. Variance isn't used very often, mainly because it is still a squared value. The measurement for variance, in this case, would be 14.67 inches *squared*, versus inches.



FORMULA TO KNOW

**Variance of a Sample**

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

**Step 5:** The final step is to take the square root of that number, which equals 3.83.

$$\sqrt{14.67} = 3.83$$

Here is an overview of the entire calculation:

$X$	$X - \bar{X}$	$(X - \bar{X})^2$
-----	---------------	-------------------

84	5.67	32.15
81	2.67	7.13
79	0.67	0.45
79	0.67	0.45
81	2.67	7.13
81	2.67	7.13
79	0.67	0.45
74	-4.33	18.75
79	0.67	0.45
71	-7.33	53.73
83	4.67	21.81
75	-3.33	11.09
81	2.67	7.13
74	-4.33	18.75
74	-4.33	18.75
Sum		205.35

$$s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$$

$$s = \sqrt{\frac{1}{15-1} [(84-78.33)^2 + (81-78.33)^2 + \dots + (74-78.33)^2]}$$

$$s = \sqrt{\frac{1}{15-1} (32.15 + 7.13 + \dots + 18.75)}$$

$$s = \sqrt{\frac{1}{15-1} (205.35)}$$

$$s = \sqrt{\frac{1}{14} (205.35)}$$

$$s = \sqrt{14.67}$$

$$s = 3.83$$

For this data set, you would expect a good portion of the heights to be within 3.83 inches of the mean, 78.33.



BIG IDEA

Variance and standard deviation share an important relationship. The variance is the square of the standard deviation, and the standard deviation is the square root of the variance.



TERM TO KNOW

#### Variance

The square of standard deviation. While it has some uses in statistics, it is not a practical unit of measurement. It is calculated the same way as standard deviation, but without the square root.

## 3. Interpreting Standard Deviation

Standard deviation is a typical amount by which we would expect values to vary around the mean.



TRY IT

Considering the original list of heights, which fall into one standard deviation of the mean, meaning their heights are either 3.83 above or below 78.33?

Height of Chicago Bulls Players		
Omer Asik	84	✗
Carlos Boozer	81	✓
Ronnie Brewer	79	✓
Jimmy Butler	79	✓
Luol Deng	81	✓
Taj Gibson	81	✓
Richard Hamilton	79	✓
Mike James	74	✗
Kyle Korver	79	✓
John Lucas III	71	✗
Joakim Noah	83	✗
Derrick Rose	75	✓
Brian Scalabrine	81	✓
Marquis Teague	74	✗
C.J. Watson	74	✗

You'll notice about  $\frac{2}{3}$  of the players had heights in this range. That is how you interpret the standard deviation.

## 4. Using Technology

As you may have noticed, calculating the standard deviation can require many steps, which could result in calculation errors. Because of this, the standard deviation is almost always found on a calculator or a spreadsheet, or some kind of applet on the internet. Typically, it is not solved by hand.

If you're frustrated with the calculation you just practiced, you can use your calculator or a spreadsheet in the future.

Here is how to use spreadsheets to calculate standard deviation:



*Step 1:* Enter your list of data into the spreadsheet.

*Step 2:* For the formula, type “=stdev(”.

*Step 3:* Highlight the list that you want to find the standard deviation for, close the parentheses, and hit "Enter."

	A	B	C	D	E
1					
2		84			
3		81			
4		79		=STDEV(B2:B16)	
5		79			
6		81			
7		81			
8		79			
9		74			
10		79			
11		71			
12		83			
13		75			
14		81			
15		74			
16		74			
17					

This spreadsheet formula finds the same number as the calculation you did by hand: 3.83.



#### BIG IDEA

If you have Excel or another spreadsheet program, there should be a standard deviation formula that you can use.



#### SUMMARY

Interpreting standard deviation is important, but it can be difficult to calculate. For this reason, calculating this number is typically done using technology. It's a measure of how far we would expect a typical data point to be from the mean. Standard deviation is the square root of a value called variance, which is a little bit easier to calculate but not as useful as a standard deviation number. Since the standard deviation is based on the mean, the standard deviation should only be reported as the measure of spread when you're reporting the measure of center to be the mean. You shouldn't mix and match the standard deviation with the median or the interquartile range.

Good luck!

Source: THIS TUTORIAL WAS AUTHORED BY JONATHAN OSTERS FOR SOPHIA LEARNING. PLEASE SEE OUR [TERMS OF USE](#).



#### TERMS TO KNOW

**Standard Deviation**



A typical amount by which we would expect a data point to differ from the mean. Typically, about half to two-thirds of the data points fall within one standard deviation of the mean.

### Variance

The square of standard deviation. While it has some uses in statistics, it is not a practical unit of measurement. It is calculated the same way as standard deviation, but without the square root.



## FORMULAS TO KNOW

### Standard Deviation of a Sample

$$s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}$$

$n$  = number of data points

$X_i$  = each of the values in the data set

$\bar{X}$  = mean of the data set

### Variance of a Sample

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$